

Executive Summary

- Diabetes mellitus (DM) and periodontal disease are chronic conditions prevalent in Hong Kong and evidence suggested that a bi-directional relationship could exist between the two diseases.
- Both DM and periodontitis can elicit inflammatory responses that mediate the damage on tooth supporting tissues and adversely affect the general health of the patients with diabetes.
- DM increases the risk and severity of periodontal disease and could compromise the response to periodontal treatment especially when glucose control is poor.
- Control of periodontal infection could potentially improve diabetic control and hence attenuate the development of diabetes complications.

Convened by:



Full English version and Chinese translation of the Position Paper on Interrelationship of Diabetes Mellitus and Periodontal Disease is available at the following websites:
Diabetes Hongkong (www.diabetes-hk.org)
Hong Kong Society of Periodontology and Implant Dentistry (www.hkspid.org)
June 2012

Interrelationship of Diabetes Mellitus and Periodontal Disease



Diabetes Mellitus and Periodontitis

Both diabetes mellitus (DM) and periodontal disease are chronic diseases prevalent in most populations.

The prevalence of Type 2 DM in Hong Kong is about 10%¹. Patients with diabetes may have no symptoms at the beginning, however, as the disease progresses, they may feel thirsty all the time, have to urinate very frequently, or have unexplained weight loss. Poor control or chronicity of DM may lead to impairment of general health and numerous complications like retinopathy, nephropathy, neuropathy, macrovascular diseases, altered wound healing and periodontitis.

Periodontitis is a chronic oral infection that results in loss of periodontal attachment, bone destruction and eventually the loss of teeth. Features of periodontal disease may include swollen and bleeding gums, gum recession, gum pocket formation, bad breath and loose teeth.

Inflammation is an important component in the pathogenesis of both diabetes and periodontitis. DM is a widely accepted major risk factor for periodontal disease and it was shown that patients with diabetes will have approximately three-fold increased risk in developing periodontal disease². A significant increase in prevalence of periodontitis has been reported in patients with type 2 DM when compared to non-diabetic controls (50% versus 36%)³. Periodontitis is therefore regarded as the sixth complication of diabetes⁴.

Conversely, periodontitis may worsen the glucose control in patients with diabetes and growing evidences suggested that the relationship between DM and periodontal disease could be bi-directional⁵.

DM Increases the Risk and Severity of Periodontal Disease

DM is a risk factor for gingivitis and periodontitis and contributes to increase in prevalence, severity and progression of periodontal disease⁶. Patients with poorly controlled diabetes may have:

1. higher risk for severe periodontal disease with odds ratio (OR) of 2.9 (for poor glucose control), OR of 1.5 (for better glucose control), when comparing with non-diabetic controls⁷.
2. more severe gingival bleeding and higher level of gingival inflammation than those with well controlled diabetes or non-diabetics controls^{8,9}
3. more advanced alveolar bone loss and periodontal destruction^{10,11}
4. worse outcome following periodontal treatment¹²
5. more recurrent periodontal infection and less favorable long term prognosis¹³

Periodontal Infection could Exacerbate the Diabetic Condition

Periodontitis has been shown to increase the risk for diabetes complications and adversely affect the glucose control in patients with diabetes¹⁴. Patients with diabetes having severe periodontitis are also more prone to diabetic complications and have a higher mortality, when compared with those without severe periodontitis¹⁵. Conversely, effective periodontal treatment appears to improve the diabetic control. Patients with type 2 DM receiving comprehensive dental treatment resulted in improvement of their glucose control with reduction of 0.9% in HbA1c when compared to those without treatment¹⁶. Provided the HbA1c reduction could be sustained, it would translate into a reduction of diabetic complications and mortality¹⁷.

Recommendations

1. Health care providers should understand the effects of diabetes on periodontal tissues and the possible impact of periodontal disease in patients with diabetes.
2. Dentists should facilitate the identification and referral of un-recognized cases of DM in dental offices. They should also control oral infection in all patients with diabetes especially those with severe periodontitis so as to reduce the inflammatory burden experienced.
3. Physicians should recognize the clinical presentations of periodontitis, enquire on its symptoms and consider oral inspection so that patients with diabetes who developed oral complications could be promptly referred to dentists for treatment.
4. Medical and dental professions should collaborate in arranging proper management of patients with diabetes in particular those with periodontitis.
5. Patients with diabetes should maintain good oral health and optimal glucose control and seek regular dental check up to minimize the adverse impact of the chronic diseases on general health.
6. Government should provide adequate resources in promoting oral health and awareness of periodontal disease in patients with diabetes; support future researches on the interrelation of both chronic diseases; and provide a platform to facilitate cross-referral between medical and dental professions for the provision of comprehensive diabetes and periodontal care.

References

1. Lam TH, Liu LJ, Janus ED, Lam KS, Hedley AJ. Fibrinogen, other cardiovascular risk factors and diabetes mellitus in Hong Kong: a community with high prevalence of Type 2 diabetes mellitus and impaired glucose tolerance. *Diabetic medicine: a journal of the British Diabetic Association*. 2000;17(11):798-806.
2. Emrich LJ, Shlossman M, Genco RJ. Periodontal disease in non-insulin-dependent diabetes mellitus. *Journal of periodontology*. 1991;62(2):123-31.
3. Leung WK, Siu SC, Chu FC, Wong KW, Jin L, Sham AS, et al. Oral health status of low-income, middle-aged to elderly Hong Kong Chinese with type 2 diabetes mellitus. *Oral health & preventive dentistry*. 2008;6(2):105-18.
4. Loe H. Periodontal disease. The sixth complication of diabetes mellitus. *Diabetes care*. 1993;16(1):329-34.
5. Taylor GW. Bidirectional interrelationships between diabetes and periodontal diseases: an epidemiologic perspective. *Annals of periodontology / the American Academy of Periodontology*. 2001;6(1):99-112.
6. Mealey BL, Oates TW. Diabetes mellitus and periodontal diseases. *Journal of periodontology*. 2006;77(8):1289-303.
7. Tsai C, Hayes C, Taylor GW. Glycemic control of type 2 diabetes and severe periodontal disease in the US adult population. *Community dentistry and oral epidemiology*. 2002;30(3):182-92.
8. Cutler CW, Machen RL, Jotwani R, Iacopino AM. Heightened gingival inflammation and attachment loss in type 2 diabetics with hyperlipidemia. *Journal of periodontology*. 1999;70(11):1313-21.
9. Ervasti T, Knuuttila M, Pohjamo L, Haukipuro K. Relation between control of diabetes and gingival bleeding. *Journal of periodontology*. 1985;56(3):154-7.
10. Salvi GE, Carollo-Bittel B, Lang NP. Effects of diabetes mellitus on periodontal and peri-implant conditions: update on associations and risks. *Journal of clinical periodontology*. 2008;35(8 Suppl):398-409.
11. Seppala B, Ainamo J. A site-by-site follow-up study on the effect of controlled versus poorly controlled insulin-dependent diabetes mellitus. *Journal of clinical periodontology*. 1994;21(3):161-5.
12. Lamster IB, Lalla E, Borgnakke WS, Taylor GW. The relationship between oral health and diabetes mellitus. *Journal of the American Dental Association*. 2008;139 Suppl:195-245.
13. Tervonen T, Karjalainen K. Periodontal disease related to diabetic status. A pilot study of the response to periodontal therapy in type 1 diabetes. *Journal of clinical periodontology*. 1997;24(7):505-10.
14. Taylor GW, Borgnakke WS. Periodontal disease: associations with diabetes, glycemic control and complications. *Oral diseases*. 2008;14(3):191-203.
15. Saremi A, Nelson RG, Tulloch-Reid M, Hanson RL, Sievers ML, Taylor GW, et al. Periodontal disease and mortality in type 2 diabetes. *Diabetes care*. 2005;28(1):27-32.
16. Lo ECM, Rong WS, Siu SC, Leung WK. Effects of dental care on glycemic control in type 2 diabetes. *The Chinese journal of dental research*. 2008;11:30-5.
17. Stratton IM, Adler AI, Neil HA, Matthews DR, Manley SE, Cull CA, et al. Association of glycaemia with macrovascular and microvascular complications of type 2 diabetes (UKPDS 35): prospective observational study. *British medical journal*. 2000;321(7258):405-12.